

AI in Perinatal Mental Health


Leveraging Artificial Intelligence for Depression and Anxiety Prediction, Diagnosis and Treatment in Low- and Middle-Income Countries

Authors

Uchechi ShirleyAnaduaka¹
Ayomide Oluwaseyi Oladosu²
Samantha Katsande³
Clinton Sekyere Frempong
Success Awuku-Amador


Affiliations:

¹University of Alberta, Canada
²The Hang Seng University of Hong Kong
³Action on Preeclampsia Ghana
School of Public Health, University of Ghana, Legon-Accra, Ghana.
Ghana Institute of Management and Public Administration




Prediction Focus

Studies using AI for risk factor identification and early detection



Treatment Focus

Studies using AI-based interventions for psychological treatment



Recent Research

Studies published since 2022, showing growing interest

Most Common Risk Factors Identified

Socioeconomic Status

Income levels and financial management capabilities consistently identified as primary risk factors.

Social Support

Family relationships and community support networks significantly impact maternal mental health.

Physical Health

BMI, previous medical conditions, and intimate partner violence are contributing factors.

Conclusions

Promising potential: AI-based methods show effectiveness in identifying risk factors for PDA

Research gap: Limited studies on AI-powered treatment interventions compared to prediction models

Ethical considerations: Many studies lacked detailed ethical frameworks for LMIC contexts

Future directions: Need for investigation into chatbot effectiveness and long-term impact assessment

Clinical and Policy Implications

Early intervention: AI tools can enhance early detection in resource-limited settings

Scalable solutions: Potential to address healthcare professional shortages

Culturally appropriate: Foundation for developing context-specific AI interventions

Policy development: Evidence base for LMIC-specific AI guidelines in maternal mental health

References

1. Gelaye, B., Rondon, M. B., Rondon, H., & Williams, M. A. (2009). Epidemiology of maternal depression, risk factors, and child outcomes in low-income and middle-income countries. *The Lancet Psychiatry*, 6(11), 935-945.

2. Obermeyer, Z., & Emanuel, E. J. (2016). Predicting the future: Big data, machine learning, and the future of medicine. *The New England Journal of Medicine*, 375(13), 1216-1219.

3. Gholizadeh, M., et al. (2020). Effectiveness of smartphone-based interventions for perinatal depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 277, 987-995.

4. Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry*, 150(6), 782-786.

5. Oh, H., et al. (2023). Artificial intelligence for mental health prediction, diagnosis, and treatment: A systematic review. *JAMA Network Open*, 6(1), e2250172.

Funding: MQ Mental Health Research Transdisciplinary Research Grant (MTGA\34)
Registration: PROSPERO CRD42024549455
Correspondence: aoladosu@hsu.edu.hk

Introduction

Perinatal depression and anxiety (PDA) affects 16-20% of women globally, with higher rates in LMICs. Despite technological advances, limited research exists on AI applications for PDA in resource-limited settings where PDA is prevalent:

- High prevalence: Up to 34% in some LMICs
- Underdiagnosed due to stigma and limited resources
- AI presents promising solutions for prediction, diagnosis, and treatment

Methods

Systematic review following PRISMA guidelines with patient and public involvement approach. Searched 7 databases for studies published January 2010-July 2024.

- PICOS framework for eligibility criteria
- Covidence software for screening and extraction
- Narrative synthesis due to study heterogeneity
- Risk assessment using Joanna Briggs Institute tools

2203

Initial Studies

Retrieved from database searches

19

Final Studies

Included after screening across 8 countries

8

Countries

China, Nepal, Brazil, India, Pakistan, Sri Lanka, Türkiye, Kenya

74.6%

Agreement Rate

Mean percentage agreement during screening

Key Results

Supervised Machine Learning Dominance

15 of 19 studies used supervised ML methods, primarily for risk factor identification and early detection rather than treatment.

Postpartum Depression Focus

14 studies (68.4%) investigated postpartum depression. Edinburgh Postpartum Depression Scale-10 was most common screening tool (42%).

Limited Treatment Applications

Only 3 conversational agents/chatbots identified for delivering psychological treatment and support.